

Antiferromagnetic Heisenberg model on anisotropic triangular lattice in the presence of magnetic field

S. Q. Shen

Department of Physics, the University of Hong Kong, Pokfulam, Hong Kong, China

F. C. Zhang

Department of Physics, University of Cincinnati, Cincinnati, Ohio 45221, USA

(Dated: July 2, 2002)

We use Schwinger boson mean field theory to study the antiferromagnetic spin-1/2 Heisenberg model on an anisotropic triangular lattice in the presence of a uniform external magnetic field. We calculate the field dependence of the spin incommensurability in the ordered spin spiral phase, and compare the results to the recent experiments in Cs_2CuCl_4 by Coldea et al. (Phys. Rev. Lett. 86, 1335 (2001)).

The ground states of two-dimensional (2D) Heisenberg models continue to be of great interest.^{1,2} In this paper, we study the antiferromagnetic (AF) Heisenberg model on an anisotropic triangular-lattice in the presence of an external magnetic field along the z-axis,

$$H_S = \frac{1}{2} \sum_{i,\delta} J_\delta \mathbf{S}_i \cdot \mathbf{S}_{i+\delta} - \mu B \sum_i \mathbf{S}_i^z. \quad (1)$$

In the above equation, the summation runs over all the lattice sites i and their neighboring sites $(i + \delta)$. We consider AF nearest neighbor spin-spin couplings, represented by J and J' as shown in Fig. 1, with both J and $J' \geq 0$. In the absence of the field, this model is equivalent to a class of models recently considered by a number of authors.^{3,4,5,6} The model includes several well known limiting cases. At $J = 0$, it is equivalent to 2D square lattice model, whose ground state is a two-sublattice Néel phase. At $J' = 0$, it becomes a set of decoupled spin chains. At $J' = J$, it is reduced to the isotropic triangular-lattice model, where the ground state is a three-sublattice antiferromagnet. Experimentally, this model may be relevant to the insulating phase of the layered molecular crystals, κ -(BEDT-TTF)₂X,⁷ and θ -(BEDT-TTF)₂RbZn(SCN)₄.⁸ Our interest on this model is largely motivated by recent experiments on Cs_2CuCl_4 . That system is a quasi 2D $S = 1/2$ frustrated Heisenberg antiferromagnet.⁹ Coldea et al.¹⁰ have used the neutron scattering to study the ground state and the dynamics of the system in high magnetic fields. Among the observations, these authors found that the incommensurate wave vector changes as the magnetic field increases, and the spiral spin density wave evolves into a fully saturated state.

In this paper we apply the Schwinger boson mean field theory (MFT) to study the effect of magnetic field in the frustrated Heisenberg models. This method enables us to study incommensurate magnetic ordering in the quantum spin systems. The magnetic ordering is identified as the Bose condensation of the Schwinger bosons, and the incommensuration of the ordering is determined by the wave vector of the condensed Schwinger bosons. In the absence of the magnetic field, the MFT pre-




FIG. 1: The anisotropic triangle lattice: the bond along the b axis is J and the bond along the a axis is J' .

dicts three possible ground state: a two-sublattice Néel phase, a spiral spin state, and a spin liquid phase, similar to the results obtained in the high temperature series expansions.³ In the presence of the field, we calculate the field dependence of the incommensuration in the spiral phase, and compare the results with the experimental observation in Cs_2CuCl_4 with good qualitative agreements.

In terms of Schwinger bosons a_{i1} and a_{i2} , the spin operators are expressed as

$$\mathbf{S}_i^+ = a_{i1}^\dagger a_{i2}; \mathbf{S}_i^- = a_{i2}^\dagger a_{i1}; \mathbf{S}_i^z = \frac{1}{2}(a_{i1}^\dagger a_{i1} - a_{i2}^\dagger a_{i2}) \quad (2)$$

with a local constraint at every site i given by $a_{i1}^\dagger a_{i1} + a_{i2}^\dagger a_{i2} = 1$. As a standard method, we introduce a Lagrangian multiplier field λ_i to describe the constraint. The Hamiltonian of the system then becomes

$$H = -\frac{1}{8} \sum_{i,\delta,\mu,\nu} J_\delta A_{ij,\mu\nu}^\dagger A_{ij,\mu\nu} - \frac{h}{2} (a_{i1}^\dagger a_{i1} - a_{i2}^\dagger a_{i2}) + \sum_i \lambda_i (a_{i1}^\dagger a_{i1} + a_{i2}^\dagger a_{i2} - 1) + \frac{1}{8} \sum_{i,\delta} J_\delta \quad (3)$$

where $h = \mu B$; $\mu, \nu = 1, 2$, $j = i + \delta$, and $A_{ij,\mu\nu} = a_{i\mu} a_{j\nu} - a_{i\nu} a_{j\mu}$ is the spin singlet operator of bond (ij) .

We note that operator $A_{ij,\mu\nu}$ is antisymmetric with respect to either the position (ij) or the indices ($\mu\nu$). On a bipartite lattice, a spin rotation by π on one of the sublattices transforms the spin-singlet bond operators into a symmetric operator with respect to the bond indices (ij). A mean field theory based on that transformation has been developed by Auerbach and Arovas.¹¹ The method was extended to study the frustrated lattices by many authors.¹² (For an overview of Schwinger boson theory, see Ref. [2] and references therein.) We introduce two types of mean fields: $\Delta_{\mu\nu}(\delta) \equiv \frac{1}{2i} \langle A_{jj+\delta,\mu\nu} \rangle$ and $\lambda = \langle \lambda_j \rangle$ where $\langle \dots \rangle$ represents the thermodynamic average. The mean field Hamiltonian may be solved using the conventional bosonic Bogliubov transformation as well as the Green function's method.

The mean field Hamiltonian in Eq. (3) is diagonalized as

$$H_{MF} = \sum_{k,\mu=\pm} \omega_\mu(k) (\alpha_{k\mu}^\dagger \alpha_{k\mu} + \frac{1}{2}) + \mathcal{E}_0, \quad (4)$$

where the single boson spectra $\omega_\mu(k) = \omega(k) \pm h/2$ with $\omega(k) = \sqrt{\lambda^2 - |\gamma(k)|^2}$ and $\gamma(k) = \sum_\delta J_\delta \Delta_{12}(\delta) \sin(k \cdot \delta)$. ($\mu = \pm$). $\alpha_{k\mu}$ is a boson annihilation operator related to the original Schwinger bosons. $\mathcal{E}_0/N_\Lambda = \sum_{\delta,\mu\nu} J_\delta (|\Delta_{\mu\nu}(\delta)|^2 + 1/4)/2 - 2\lambda$. The free energy is given by ($\beta = 1/k_B T$, with T the temperature)

$$F = \frac{1}{\beta} \sum_{k,\mu=\pm} \ln[1 - \exp[-\beta\omega_\mu(k)]] + \frac{1}{2} \sum_{k,\mu=\pm} \omega_\mu(k) + \mathcal{E}_0. \quad (5)$$

The mean field Hamiltonian is solved together with the self-consistent equations for the two types of mean fields, which are given by

$$\frac{1}{N_\Lambda} \sum_{k,\mu=\pm} \frac{\lambda}{\omega(k)} [n_B(\omega_\mu(k)) + 1/2] = 2; \quad (6a)$$

$$\frac{1}{N_\Lambda} \sum_{k,\mu=\pm} \frac{\gamma(k)}{\omega(k)} \sin(k \cdot \delta) [n_B(\omega_\mu(k)) + 1/2] = 2\Delta(\delta). \quad (6b)$$

In the MFT, the magnetic ordering may be identified as the Schwinger boson condensation.¹³ Below we examine the possible Schwinger boson condensation at the wave vector $k = \pm k^*$, corresponding to the lowest energy of $\tilde{\omega}_\mu(\pm k^*) \rightarrow 0$. We introduce a non-negative quantity,

$$b_0 = \frac{2\lambda}{N_\Lambda \omega_-(k^*)} [n_B(\omega_-(+k^*)) + n_B(\omega_-(-k^*))], \quad (7)$$

such that the points $k = \pm k^*$ in the integrals in Eqs.(6a) and (6b) are taken into account separately,

$$\sum_{\mu=\pm} \int \frac{dk'}{(2\pi)^2} \frac{\lambda}{\omega(k')} [n_B(\omega_\mu(k')) + 1/2] = 2 - b_0; \quad (8a)$$

$$\sum_{\mu=\pm} \int \frac{dk'}{(2\pi)^2} \frac{\gamma(k')}{\omega(k')} \sin(k' \cdot \delta) [n_B(\omega_\mu(k')) + 1/2] = \frac{2J_\delta \Delta(\delta)}{J_\delta} - b_0 \gamma(k^*) \sin(k^* \cdot \delta)/\lambda. \quad (8b)$$

The general features of the mean field solutions in 2D lattices are qualitatively given as below. At any finite T , $b_0 = 0$. At $T = 0$, if $\min[\omega(k)] \neq 0$ and $b_0 = 0$, the ground state is a spin liquid, whose gap depends on the minimal value of the spectra $\omega(k)$. If $\min[\omega(k)] = 0$ and b_0 is finite, the Schwinger bosons are condensed to the lowest energy state, and the system possesses the magnetic long-range order. The ordering wave vector Q is determined by $\omega(k^*) = 0$. From the spin-spin correlations,

$$\chi_{\alpha\alpha}(q) = - \lim_{\tau \rightarrow 0^-} \langle T_\tau \mathbf{S}_q^\alpha(\tau) \mathbf{S}_{-q}^\alpha(0) \rangle \quad (\alpha=x,y,z),$$

we have

$$\begin{aligned} \chi_{xx}(q) &= \chi_{yy}(q) = \chi_{zz}(q) \\ &= \frac{1}{4N_\Lambda} \sum_k \left(\frac{n_B[\omega(k)] + \frac{1}{2}}{\omega(k)} \right) \left(\frac{n_B[\omega(k+q)] + \frac{1}{2}}{\omega(k+q)} \right) \\ &\quad \times [\lambda^2 - \gamma(k)\gamma(k+q)] - \frac{1}{16} \end{aligned} \quad (9)$$

for $h = 0$, which indicates the mean field theory does not break SU(2) symmetry. In the thermodynamic limit, ($N_\Lambda \rightarrow +\infty$), the correlation functions are convergent except for $Q = 2k^*$,

$$\frac{\chi_{\alpha\alpha}(Q)}{N_\Lambda} = \frac{b_0^2}{8} \quad (10)$$

which indicates that there exists long-range correlations with $Q = 2k^*$ whence $b_0 \neq 0$.

On the triangular-lattice (see Fig. 1), each site has six neighbors: $\pm \mathbf{a}_1, \pm \mathbf{a}_2, \pm(\mathbf{a}_1 + \mathbf{a}_2)$. It is convenient to write the wave vector $k = (k_1, k_2)$, with k_1 and k_2 the components of the vector along the directions of a_1 and a_2 , respectively. The lattice constant a is set to be 1.

We now consider the mean field solutions at $T = 0$. Let me first discuss the solutions in the absence of the field. At $J'/J = 1$ (the isotropic triangular lattice), $k_0^* = \pi/3$ and $\omega(k^*) = 0$, b_0 is finite indicating a magnetic long range order with the ordering wave vector $Q(2\pi/3, 2\pi/3)$. At $J = 0$ (the square lattice), $k_0^* = \pi/2$ and $\omega(k^*) = 0$, and $b_0 > 0$ implying a Neel ordering at $Q(\pi, \pi)$. The MFT in these two limiting cases is consistent with the known results. At $J' = 0$ (decoupled one-dimensional chains), the MFT gives $k_0^* = \pi/4$, ($2k^*$ along the chains),




FIG. 2: The critical field in unit J versus the ratio J'/J .

$\omega(k^*) > 0$, and $b_0 = 0$, suggesting a spin gap state. The 1D model is exactly solvable, and the ground state is a gapless spin liquid¹⁴, although the static spin correlation becomes strongest at $Q_b = \pi$.¹⁵ The discrepancy between the MFT and the exact solutions is primarily due to the neglect of the topological term in the MFT. For the general values of J'/J , the MFT predicts three phases at $h = 0$: (1). a spin liquid phase at $J'/J < 0.136$; (2). a spin spiral state at $0.136 < J'/J < 1.70$, with the ordering wave vector between $(\pi/2, \pi/2)$ and (π, π) ; (3). an antiferromagnet with the ordering wave vector $Q = (\pi, \pi)$ at $J'/J > 1.70$. We note that the phase diagram of the model was studied previously by using a series expansion technique, linear spin density wave and SP(N) mean field theory etc.^{3,5,6,12} In the method of series expansion, they found a Néel state persisting down to $J'/J > 1.43$, and predicts a spiral phase at small ratio of J'/J . These are consistent with our MFT. These authors also found a dimer phase between the Néel and spiral states. The dimer phase breaks translational invariance and is not included in the present MFT.

We now consider the solutions at $h \neq 0$. In this case, the SU(2) rotational symmetry is broken, and $\omega_-(k) < \omega_+(k)$. The Bose condensation criterion is given by $\omega_-(k^*) = \omega(k^*) - h/2 = 0$. We use the MFT to study the field dependent incommensurability in the spin spiral phase. The results are shown in Fig. 2, where the vector k^* are plotted as a function of J'/J at several values of h . The main feature is as follows. 1) $J'/J = 1$ is a stable fixed point, around which the external magnetic field does not change the ordering wave vector; 2) At $0.136 < J'/J < 1$, the ordering wave vector increases slightly as the field increases; 3) At $J'/J > 1$, the ordering wave vector decreases as the field increases; 4) At $J'/J < 0.136$, the spin liquid may evolve into a spiral state, and becomes fully saturated as the field further increases.

In the presence of the magnetic field, the spin z -component $\langle S_z \rangle \neq 0$, and the ground state breaks the SU(2) invariance. At $T = 0$, the expectation value of




FIG. 3: The ratio versus the wave vector k^* at different fields are plotted. Inset: The wave vector near $J'/J = 0.33$ at ten different fields are plotted. From left to right, $h^* = 0.0, 0.1, \dots, 1.0$.

S_z is given by

$$\langle S_i^z \rangle = \frac{1}{2} h^* b_0(k^*, h^*), \quad (11)$$

where $h^* = h/(2\lambda)$ is the dimensionless field. At $h \neq 0$, $\langle S_i^z \rangle$ is finite, which indicates a polarized component along the field orientation z -axis. The static transverse susceptibilities at $Q = 2k^*$ are given by

$$\frac{\chi_{\alpha\alpha}(Q)}{N_\Lambda} = \frac{b_0^2(k^*, h^*)}{8} [1 - (h^*)^2] \quad (12)$$

for $\alpha = x$ and y . $\chi_{\alpha\alpha}(Q)/N_\Lambda$ decreases as the field increases, and approaches to zero at $h^* = 1$. $b_0(k^*, h^*)$ can be calculated within the MFT. A special case is $b_0(k^*, h^* = 1) = 1$ corresponding to $\langle S_i^z \rangle = 1/2$, or the spin full polarization. We have calculated the critical field μB_c (defined as the lowest field to induce the full spin polarization) as a function of J'/J . The results are plotted in Fig. 3. As we can see, μB_c increases as J' increases for the fixed J . As J' increases, the spin couplings are strengthened, and it requires a higher field to polarize the spin.

Very recently, Coldea et al.¹⁰ have reported the neutron scattering experiments on the antiferromagnet Cs_2CuCl_4 in the high magnetic field. That system is a quasi-2D spin-1/2 quantum system in a triangular-lattice as shown in Fig.1. The spin-spin couplings are anisotropic with $J'/J \approx 0.33$. In the absence of the external magnetic field, the spins are incommensurately ordered and are aligned within the plane of the triangular-lattice. The latter may indicate a weak deviation from the Heisenberg model. Coldea et al. have studied the low temperature states of the system in the presence of in-plane as well as perpendicular magnetic fields. In the presence of the perpendicular fields, the states are found magnetically ordered with the varying incommensuration below a critical field, above which the system becomes fully spin polarized ferromagnet. In the presence of the




FIG. 4: The incommensuration ϵ versus the external field at $J'/J = 0.33$.

in-plane field, they have observed additional spin liquid phase between the incommensurate states and the ferromagnetic phase. There have been theoretical efforts to understand their experimental results.¹⁶ In the present paper, we have only considered the Heisenberg model in a uniform magnetic field. The predicted spin structure breaks SU(2) symmetry, and shows the spin polarization along the field-direction. Such a spin structure is compatible with the experiments in the perpendicular field, but incompatible with the in-plane field. Therefore, our MFT may be of relevance to the perpendicular field case in their experiments.

To compare with the experiments, we define a quantity to describe the incommensuration, which is proportional to the wave vector deviation from the Néel state, $\epsilon = 1/2 - Q_1/\pi$, with Q_1 the component of the spiral wave vector along a_1 direction. In Table I, we list the experimental¹⁰ and theoretical values of ϵ at field $h^* = 0$ and at the critical field $h^* = 1$ for the two values of J'/J . Also listed are the values of ϵ at $h^* = 0$ calculated from

the series expansion method³. The agreements between the present MFT and the series expansion at $h^* = 0$ are very good. In Fig. 4, we plot the incommensuration relative to the Néel state as a function of the external field for $J'/J = 0.33$. Our mean field results are in qualitative agreement with the experiments: as field increases, ϵ also increases in the parameter space of interest. Quantitatively, the theory predicts a weaker variation in the incommensuration than in the experiments. This discrepancy could be partly due to the neglect of the deviation of the physical system from the Heisenberg model in the theory.

h^*	Exp. ¹⁰	$J'/J = 0.404$	$J'/J = 0.33$	SE ³
0.0 (ϵ_0)	0.03	0.047	0.035	0.028
1.0 (ϵ_c)	0.053	0.053	0.0422	—

Table I: The incommensuration to the Néel state along the b axis is listed. $h^* = 0$ means the absence the magnetic field, and $h^* = 1$ means the state is saturated fully. SE means the series expansion method, and the data is estimated from the work by Zheng et al.³, which was also predicted by Manuel and Ceccatto.⁵

In summary, we have used a Schwinger boson mean field theory to study the antiferromagnetic Heisenberg model on an anisotropic triangular lattice in the presence of an external magnetic field. We calculate the magnetic field dependence of the incommensurability of the spin spiral phase. The theoretical results are compared well with the recent neutron scattering experiments.

We would like to thank stimulating discussions with D. A. Tennant and R. Coldea on their experiments. This work was supported by a RGC grant of Hong Kong and a CRCG grant of the University of Hong Kong, and by the US DOE grant No. FG03-01ER45687, and by the Chinese Academy of Sciences. The authors also acknowledge ICTP at Trieste, Italy for its support and hospitality, where part of the work was initiated.

¹ S. Chakravarty, B. I. Halperin, and D. R. Nelson, Phys. Rev. B 39, 2344 (1989).

² A. Auerbach, Interacting Electrons and Quantum Magnetism (Springer-Verlag, New York, 1994).

³ Zheng Weihong, R. H. McKenzie, and R. R. P. Singh, Phys. Rev. B 59, 14367 (1999).

⁴ J. Merino, R. H. McKenzie, J. B. Marston, and C. H. Chung, J. Phys.: Condens. Matter 11, 2965 (1999).

⁵ L. O. Manuel and H. A. Ceccatto, Phys. Rev. B 60, 9489 (1999).

⁶ C. H. Chung, J. B. Marston, and R. H. McKenzie, J. Phys. Condens. Mater. 13, 5159 (2001).

⁷ R. H. McKenzie, Comments Condens. Matter Phys. 18, 309 (1998).

⁸ J. Schmalian, Phys. Rev. Lett. 81, 4232 (1998); H. Kino and H. Kontani, J. Phys. Soc. Jpn. 67, 3691 (1998); H. Kino and T. Moriya, J. Phys. Soc. Jpn. 67, 3695 (1998); M. Votja and E. Dagotta, Phys. Rev. B 59, 713 (1999).

⁹ R. Coldea, D. A. Tennant, R. A. Cowley, D. F. McMorrow, B. Dorner, and Z. Tylczynski, Phys. Rev. Lett. 79, 151 (1997).

¹⁰ R. Coldea, D. A. Tennant, A. M. Tsvelik, and Z. Tylczynski, Phys. Rev. Lett. 86, 1335 (2001).

¹¹ A. Auerbach and D. P. Arovas, Phys. Rev. Lett. 61, 617 (1988); D. P. Arovas and A. Auerbach, Phys. Rev. B 38, 316(1988).

¹² I. Richey and P. Coleman, J. Phys. Condens. Mater. 2, 9227 (1990); D. Yoshioka and J. Miyazaki, J. Phys. Soc. Jpn. 60, 614 (1991); A. Mattsson, Phys. Rev. B 51, 11574 (1995).

¹³ S. Sarker, C. Jayaprakash, H. R. Krishnamurthy, M. Ma, Phys. Rev. B 40, 5028 (1989).

¹⁴ H. Bethe, Z. Phys. 71, 205 (1931).

¹⁵ S. Q. Shen, Inter. J. Mod. Phys. 12, 709 (1998).

¹⁶ M. Bocquet et al., Phys. Rev. B 64, 094425 (2001).